






DISTINCTION



Need scoops in quarters.


Bruising
pink

Right Front



High inside


Left front




Pink
bruising

under run heels

Right Hind



Left Hind



too long - breakover
is here back

very high side



DISTINCTION

Equine Evaluation Form

Age	12
Breed	QH
Primary discipline	All Around/heavy workload, trail, arena, ranch work

Current Body State

Natural Asymmetry / Crookedness pattern	Neck contracted right , barrel contracted left/ dominant right fore
Hoof condition/balance	Barefoot. Experiencing significant (sore foot) lameness in all 4 feet. Irregular hairline/'jamming upward' coronet bands above quarters. Balance, OK, but recommend scoops in quarters and appropriate beveling of edges. Overall reduction in protein in the diet (Free feed grass hay, (Teff/Timothy combo or Bermuda /Timothy) Bucket daily consisting of: loose beet pulp shreds (2 lbs dry measure) soaked, along with 2 tablespoons Larch powder. Access to free choice: loose white salt, ABC-GRP, A-Mix, & BVC-Mix
Assessment Notes:	Overall 'deflated' appearance. Ulcer/stress dips in skull above eyes. Experiencing common symptoms of crookedness. Heavy in front, very stuck in front end, particularly on right front. "U" forming in neck in front of withers, Sternum dropping through the chest. Back taking on suspension bridge appearance. Left side: Inappropriate bulging of the splenius muscle (upper neck region), shear line along scapular spine, and another along the deltoid muscle and diffuse tendon of origin (angle along slope of shoulder) reactive to all ulcer points, tight in lumbar sacral junction and down lumbodorsal fascia. Left stifle sore, slow reflex response, left hock feels hot. Right side: Right jaw more pronounced, right ear slightly higher. Same bulge in neck and shear lines along shoulder, additional shear line in the cervical part of the ventral serrate muscle on left side of the neck. Saddle scars behind withers, more significant on right side than left. Sore in lumbar region (right loin) and Iliacus muscle (lower right hip region)

Behavior manifestations/patterns	Willing & kind of horse, full of try. However, he is in a state near total postural collapse. Experiencing difficulty moving forward with any engagement of a postural ring. Disconnected from hind end to front end as well as front end to hind end.
Recommendations/Training:	30 days groundwork: focusing on proper bending, contact yielding, connecting front end to hind/ hind end to front end. Caveson work in-hand to strengthen, rebalance, and begin straightening and shifting center of mass out of shoulders, back. Actively engage hindlegs through diagonals to begin to equalize and develop more carry, less push. Followed by 30 days continuing caveson work, and reintroduction of ridden work, SLOWLY (following classical principles & gymnastic riding). *Suggesting hoofboots for hoofcare transition. All links for recommended supplements, boots, etc. provided in email body.